arXiv:solv-int/9710008v2 7 Nov 1997

On classical string configurations

B.G. Konopelchenko and G. Landolfi
Dipartimento di Fisica, Universitá di Lecce, 73100 Lecce, Italy
and
I.N.F.N., Sezione di Lecce, 73100 Lecce, Italy

February 9, 2008

Abstract

Equations which define classical configurations of strings in R^3 are presented in a simple form. General properties as well as particular classes of solutions of these equations are considered.

Polyakov's theory of strings [1]-[2] has been discussed in a number of papers during the last ten years (see *e.g.* [3]- [4]). Various problems and features of this theory have been studied and analyzed. Stable classical configurations of strings is one of the important problems of the string theory. Their description, classification and analysis of their properties are essential for understanding classical, quasi-classical and other aspects of the string theory. Surprisingly, not many results concerning classical configurations of strings (world-sheets) have been obtained (see *e.g.* [5]-[11]). The reason, obviously, lies in the complexity of the corresponding equations. So the description and analysis of the classical configurations of string is still of a great interest.

In the present paper, we propose a new approach to the problem of classical configurations of strings in the three-dimensional Euclidean space. It is based on the generalized Weierstrass formulae which allows to construct any surface in R^3 starting with a system of two linear equations. Then we represent the Euler-Lagrange equation for the Nambu-Goto-Polyakov action in a simple form. Common solutions of this equation and above linear system provide surfaces in R^3 which describe classical configurations of strings (world-sheets). We discuss both general properties of the above system of equations as well as particular classes of solutions.

Generalized Weierstrass formulae proposed in [12] are the extension of well-known formulae of Weierstrass for minimal surfaces. One starts with the following system of two linear equations [12]:

$$\begin{cases} \psi_{1z} = p\psi_2 \\ \psi_{2\overline{z}} = -p\psi_1 \end{cases} \tag{1}$$

where $\psi_1(z,\overline{z}), \psi_2(z,\overline{z})$ are complex-valued functions, $p(z,\overline{z})$ is a real-valued function and $z,\overline{z}$ are complex variables. Then one defines three real-valued functions X_1, X_2, X_3 as follows [12]:

$$\begin{aligned} X_1 &+ iX_2 = 2i\int_\Gamma \left(\overline{\psi}_1^2 dz' - \overline{\psi}_2^2 d\overline{z}'\right) \\ X_1 &- iX_2 = 2i\int_\Gamma \left(\psi_2^2 dz' - \psi_1^2 d\overline{z}'\right) \\ X_3 &= -2\int_\Gamma \left(\psi_2\overline{\psi}_1 dz' + \psi_1\overline{\psi}_2 d\overline{z}'\right) \end{aligned} \tag{2}$$

where Γ is an arbitrary path of integration in the complex plane and bar means complex conjugation. Due to (1), the integrals in (2) do not depend on the choice of Γ. Finally, one treats X_1, X_2, X_3 as the coordinates of a surface in R^3 [12]. The formulae (2) define the conformal immersion of a surface in R^3 since the corresponding first fundamental form is

$$d\ell^2 = 4u^2 dz d\overline{z} \tag{3}$$

where $u = |\psi_1|^2 + |\psi_2|^2$. From (3) it is clear that $z,\overline{z}$ play the role of minimal lines ($z = x+iy, \overline{z} = x - iy$). The Gaussian curvature K and the mean curvature H look like:

$$K = -\frac{1}{u^2}\left(\log u\right)_{z\overline{z}} \quad , \quad H = \frac{p}{u} \tag{4}$$

The generalized Weierstrass formulae (1)-(2) are the powerful tool for studying surfaces in R^3. In particular, any analytic surface in R^3 can be globally represented by the formulae (1)-(2) [13]. Note that another extension of the Weierstrass formulae for non-minimal surfaces has been proposed earlier in [14]. The equivalence of the generalized Weierstrass formulae (1) and those of [14] has been established in [15]. However the form (1) is more convenient for many purposes.

The generalized Weierstrass formulae are also useful for an analysis of various quantities and problems of Polyakov string theory. Let us consider the standard Nambu-Goto-Polyakov action [1]-[2]

$$S = \mu_0 \int d\Sigma + \frac{1}{\alpha_0} \int H^2 d\Sigma \tag{5}$$

where $d\Sigma$ is the area element. In terms of quantities p, ψ_1, ψ_2, from (1)-(2) it has the following form [15]:

$$S = 4\mu_0 \int \left(|\psi_1|^2 + |\psi_2|^2\right)^2 dxdy + \frac{4}{\alpha_0} \int p^2 dxdy \tag{6}$$

In the infrared region, where the second term in (5) dominates [2], the action S becomes quadratic in p and the generating Polyakov's integral over all surfaces is reduced to a very simple one:

$$Z \sim \int [d\sigma]\, e^{-\frac{4}{\alpha_0}\int p^2 dxdy} \tag{7}$$

The measure of integration $[d\sigma]$ depends in general both on p and ψ_1, ψ_2. We will discuss the functional (7) in a separate paper. Here, we will consider classical configurations of strings which are given by solutions of the Euler-Lagrange equation for the action (5).

In generic coordinates the corresponding Euler-Lagrange equation has the form [16]-[17]:

$$\Delta H + 2H\left(H^2 - K\right) - 2\alpha_0\mu_0 H = 0 \tag{8}$$

where Δ is the Laplace-Beltrami operator. If one chooses on a surface a conformal metric as in (3), then

$$\Delta H = \frac{1}{u^2} H_{z\bar{z}} \tag{9}$$

and in terms of the variables $\varphi = H^{-1}$ and $p = u/\varphi$ equation (8) takes the form:

$$\varphi_{z\bar{z}} + \left[2p^2 + \left(\log p^2\right)_{z\bar{z}}\right]\varphi - 2\alpha_0\mu_0 p^2 \varphi^3 = 0 \tag{10}$$

Note that the Euler-Lagrange equation (8) has the form (10) in the conformal metric independently of any use of the generalized Weierstrass formulae (1)-(2). But if one starts with the Weierstrass formulae (1)-(2), then the quantity p in (10) is exactly the coefficient p in equation (1) and $\varphi = \left(|\psi_1|^2 + |\psi_2|^2\right)/p$. Thus equations (1), (10) form a system of equations which completely defines the classical configurations of strings in R^3 via the formulae (2). Any surface in R^3 is constructable by the formulae (1)-(2). Those which obey in addition equation (10) represent classical configurations of strings. The system of equation (1), (10) is much simpler than those in terms of coordinates X_1, X_2, X_3 derived and used in [5]- [9]. Equation (10) itself (without the generalized Weierstrass formulae (1)-(2)) defines possible candidates to the classical configurations of strings. So it is of the great interest and importance itself. We will present below some classes of its solutions.

We will consider here three particular cases: *1)* constant mean curvature; *2)* constant mean curvature density; *3)* the case $\mu_0 = 0$. The simplest solution of equation (10) corresponds to the constant mean curvature, *i.e.* to $\varphi = \varphi_0 = \text{constant}$. In this case equation (10) is reduced to the Liouville equation for $\theta = 2\log p$:

$$\theta_{z\bar{z}} + \beta e^{\theta} = 0 \tag{11}$$

where $\beta = 2\varphi_0 - 2\alpha_0\mu_0\varphi_0^3$. The general solution of equation (11) is of the form (see *e.g.* [18])

$$\exp\theta = \frac{A_z \overline{A}_{\bar{z}}}{\left(|A|^2 + \beta/2\right)^2} = p^2 \tag{12}$$

while for $\beta = 0$ one has $p^2 = A(z)\overline{A}(\bar{z})$ where $A(z)$ is an arbitrary analytic function. So the formula (12) gives us two cylindrical surfaces ($\beta = 0$) with $H_0 = \pm\sqrt{\alpha_0\mu_0}$. At $H > \sqrt{\alpha_0\mu_0}$ and $-\sqrt{\alpha_0\mu_0} < H < 0$ surfaces are spherical ($\beta > 0$ and, hence, $K > 0$) while for $H < -\sqrt{\alpha_0\mu_0}$ and $0 < H < \sqrt{\alpha_0\mu_0}$ one has pseudo-spherical surfaces ($\beta < 0$ and $K < 0$).

Now we consider as second particular case $H\sqrt{g} = const.$, namely $p = p_0 = const.$ Equation (10) is now

$$\varphi_{z\bar{z}} + 2p_0^2\varphi - 2\alpha_0\mu_0 p_0^2\varphi^3 = 0 \tag{13}$$

In the simplest case $\mu_0 = 0$, we have the linear equation

$$\varphi_{xx} + \varphi_{yy} + 8p_0^2\varphi = 0 \tag{14}$$

General solution and classes of particular solutions of this equation are easily available. For the square domain $0 \leq x, y \leq \pi$ and vanishing φ at the boundary, one has the well-known family of solutions

$$\varphi_{nm} = \frac{1}{H_{nm}} = A_{nm} \sin nx \ \sin my \tag{15}$$

where $n, m = 1, 2, \ldots$, A_{nm} are constants and p_0 takes discrete values $p_0^2 = (n^2 + m^2)/8$. So the corresponding closed surface has the following first fundamental form, string action and total area

$$\begin{aligned} d\ell^2 &= \frac{n^2+m^2}{2} \left(A_{nm} \sin nx \ \sin mx\right)^2 \left(dx^2 + dy^2\right) && (16) \\ S_{nm} &= \frac{(n^2+m^2)}{2\alpha_0} && (17) \\ \Sigma_{nm} &= \frac{\pi^2 (n^2+m^2) A_{nm}^2}{2} && (18) \end{aligned}$$

respectively. Equation (13) is well-known in various fields of physics. It has the famous kink-antikink solution [19]:

$$\varphi = \pm \left(\alpha_0 \mu_0\right)^{-1/2} \tanh \left[p_0 \left(e^{ia} z + e^{-ia} \overline{z} + b\right)\right] \tag{19}$$

where a, b are arbitrary real constants. It must be noted that the mean curvature $H = \varphi^{-1}$ is singular along the line $2x \cos a - 2y \sin a + b = 0$. Let us consider now the case when φ depends only on $x = (z + \overline{z})/2$. Equation (13) is equivalent to

$$\varphi_x^2 + 8p_0^2 \varphi^2 - 4\alpha_0 \mu_0 p_0^2 \varphi^4 = const. \tag{20}$$

Solution of this equation are given by standard elliptic functions. Equation (13) arises also in a more general case, namely when $(\log p)_{z\overline{z}} = 0$. Since then $p^2 = A(z)\overline{A}(\overline{z})$ -where $A(z)$ is an arbitrary analytic function- equation (10) is reduced to (13) after the change of variable $z \to A(z)$.

Among the case when both $p \neq const$ and $\varphi \neq const$ we mention one with $p^2 = 2A_z \overline{A}_{\overline{z}} \left(A + \overline{A}\right)^{-2}$ where $A(z)$ is an arbitrary analytic function. Changing $z \to \xi = A(z)$, one gets from (10) the equation

$$(\xi + \overline{\xi})^2 \varphi_{\xi\overline{\xi}} + 6\varphi - 4\alpha_0 \mu_0 \varphi^3 = 0 \tag{21}$$

In the pure Polyakov case $\mu_0 = 0$ this equation becomes linear and general solution is of the form (see [20]) $\varphi = \left(\partial_\xi - \frac{3}{\xi+\overline{\xi}}\right)^2 B(\xi) + c.c.$ where $B(\xi)$ is an arbitrary analytic function.

Among the different possible constraints on p and φ there is one of importance in string theory. It is the gauge of constant mean density $H\sqrt{g} = 1$ in which the Virasoro symmetry is easily revealed (see *e.g.* [3]). In our variables it is the constraint $p^2\varphi = 1$. In this gauge one has $S = 4\int \left(\mu_0\varphi + \alpha_0^{-1}\varphi^{-1}\right) dxdy$ and equation (10) looks like

$$\varphi_z\varphi_{\overline{z}} + 2\varphi - 2\alpha_0\mu_0\varphi^3 = 0 \tag{22}$$

In the one-dimensional case (say $\varphi_y = 0$), setting $\varphi = (2\alpha_0\mu_0)^{-1}\wp$ this equation is reduced to

$$\wp_x^2 = 4\wp^3 - 16\alpha_0\mu_0\wp \tag{23}$$

that is the standard equation for the Weierstrass elliptic function with the invariants $g_2 = 16\alpha_0\mu_0$ and $g_3 = 0$. Equation (22) is also solvable by the elliptic change of the dependent variable and then by the use of the method of characteristic. For example, this procedure gives

$$\varphi = \wp\left(c + \sqrt{2\alpha_0\mu_0}\sqrt{(x-a)^2 + (y-b)^2}\right) \tag{24}$$

where c, a and b are constants and now $\wp$ is the Weierstrass elliptic function with the invariants $g_2 = 4/\alpha_0\mu_0$ and $g_3 = 0$.

Now we will consider our third particular case $\mu_0 = 0$, when equation (10) is reduced to the linear equation

$$\varphi_{z\overline{z}} + V(z,\overline{z})\varphi = 0 \tag{25}$$

where $V = 2\left[(\log p)_{z\overline{z}} + p^2\right]$. For periodic functions p we have $\int H^2 d\Sigma = 4\int H^2 dxdy$. So the action (5) in this case is completely defined by the *potential* $V(z,\overline{z})$. If $V = const$, we have an infinite family of solutions for equation (25). To define p one has to solve the equation $\theta_{z\overline{z}} + 2\exp\theta = V = const.$, where $\theta = p^2$. The existence of a wide class of periodic solutions of this equation was claimed in [22]. Using those results and solution of equation (14) presented above, one can construct the variety of closed surfaces with the same value of action $S = 2\pi^2 V/\alpha_0$.

In the one-dimensional case $\varphi = \varphi(x)$ one can use the known results about the one-dimensional Schrödinger equation $-\varphi_{xx}+Q(x)\varphi = E\varphi$. There are several solvable cases. One of them is provided by the potential $Q = \frac{2}{\sin x^2}$ (Sutherland model [24]). Taking the periodic wave function $\varphi = \frac{2\sqrt{2}}{\sin x}$ associated with the unphysical value of the energy $E = 1$, one has $V = \frac{1}{4} - \frac{1}{2\sin x^2}$. Calculating p^2, one gets $p = \frac{\sin x}{2\sqrt{2}\left(\sqrt{2}-\sin x\right)}$. Thus $H = \frac{\sin x}{2\sqrt{2}}$ and $u = \left(\sqrt{2} - \sin x\right)^{-2}$. The corresponding surface is nothing but the Clifford torus (see [16]). Here we would like to emphasize its connection with the solvable (Sutherland) case of equation (25). Other solvable potentials in (25) may provide remarkable surfaces too. Nevertheless, it must be pointed out that the requirement of a physical interpretation is crucial and we must not forget that we are dealing with real surfaces. For instance, for $Q = -\frac{2}{\cosh x^2}$ (Bargmann potential) and $E = 1$, one has a real mean curvature $H = \frac{\cosh x}{2\sqrt{2}}$ and pure imaginary $p = \frac{i\cosh x}{2\sqrt{2}\left(\sqrt{2}+\cosh x\right)}$. Consequently, the metric is negatively defined: $u^2 = -\left(\sqrt{2} + \cosh x\right)^{-2}$. Here, the reality of p is the problem.

A variety of exact solutions for the two-dimensional Schrödinger equation and, hence, equation (25) is available too. They have been constructed by the inverse-spectral transform method (see *e.g.* the review [25]) and are expressed in terms of the Prym θ-function.

Now we will discuss some examples of solutions of system of equations (1), (10) and the corresponding surfaces given by formulae (1)-(2). The first example is given by

$$p = \frac{1}{4}, \quad \psi_1 = \psi_2 = \frac{\sqrt{R}}{2} \exp \frac{iy}{2} \tag{26}$$

So the potential $V = 1/8$ is constant and we get a cylinder of radius R:

$$X_1 = -R\sin y, \quad X_2 = -R\cos y, \quad X_3 = -Rx \tag{27}$$

The second case is

$$\begin{aligned} p &= 0 \\ \psi_1 &= \frac{1}{2}\exp\left(\frac{iy-x}{2}\right), \quad \psi_2 = \frac{1}{2}\exp\frac{iy+x}{2} \end{aligned} \tag{28}$$

and

$$X_1 = -\sin y \ \cosh x, \quad X_2 = -\cos y \ \cosh x, \quad X_3 = -x \tag{29}$$

namely the catenoid. The third case is

$$\begin{aligned} p &= \frac{1}{2\cosh x} \\ \psi_1 &= \frac{1}{2\cosh x}\exp\left(\frac{iy+x}{2}\right), \quad \psi_2 = \frac{1}{2\cosh x}\exp\frac{iy-x}{2} \end{aligned} \tag{30}$$

and

$$X_1 = -\frac{\sin y}{\cosh x}, \quad X_2 = -\frac{\cos y}{\cosh x}, \quad X_3 = -\tanh x \tag{31}$$

that is the unit sphere $X_1^2 + X_2^2 + X_3^2 = 1$. Also in this case $V = 0$.

Fourth example : the Clifford torus discussed above (which can be obtained by rotation of the circle of radius 1 with the centre at $(\sqrt{2}, 0)$ corresponds to

$$\begin{aligned} p &= \frac{\sin x}{2\sqrt{2}\left(\sqrt{2} - \sin x\right)} \\ \psi_1 &= \frac{\sqrt{\left(\sqrt{2} - \sin x - \cos x\right)}}{2\left(\sqrt{2} - \sin x\right)} e^{\frac{iy}{2}}, \quad \psi_2 = \frac{\sqrt{\left(\sqrt{2} - \sin x + \cos x\right)}}{2\left(\sqrt{2} - \sin x\right)} e^{\frac{iy}{2}} \end{aligned}$$

and

$$X_1 = -\frac{\sin y}{\sqrt{2} - \sin x}, \quad X_2 = -\frac{\cos y}{\sqrt{2} - \sin x}, \quad X_3 = -\frac{\cos x}{\sqrt{2} - \sin x} \tag{32}$$

The potential is $V = \frac{1}{4} - \frac{1}{2\sin^2 x}$. For the Clifford torus $\int H^2 d\Sigma = 2\pi^2$. According to the conjecture of Wilmore [16], the Clifford torus provides the minimum for the functional $\int H^2 d\Sigma$ within a class of compact surfaces of genus 1.

And finally, the fifth example which is associated to the potential

$$V(x) = \frac{R^4 - 8R(R - \sin(f))}{8[R - 2\sin(f)]^2}$$

Here $f_x(x) = R - \sin[f(x)]$, *i.e.*

$$f(x) = 2\arctan h(x)$$

with $h(x) = \frac{1}{R} + \sqrt{1 - \frac{1}{R^2}} \tan\left[\sqrt{R^2-1}(x - C_1)/2\right]$ from which follows that

$$\sin[f(x)] = \frac{2h(x)}{1 + h^2(x)}$$

The corresponding surface, defined by

$$p = \frac{R - 2\sin[f(x)]}{4}, \quad \psi_1 = \sqrt{\frac{f - f_x}{4}} e^{\frac{iy}{2}}, \quad \psi_2 = \sqrt{\frac{f + f_x}{4}} e^{\frac{iy}{2}} \tag{33}$$

and, consequently,

$$X_1 = -f(x)\cos y, \quad X_2 = -f(x)\sin y, \quad X_3 = -\cos f(x) \tag{34}$$

represents an extension of the Clifford torus [26].

Other analytic and numerical solution of equations (1)-(10) and corresponding classical configurations of strings will be discussed elsewhere.

REFERENCES

1. A.M. Polyakov, Phys. Lett. **B103**, 207, (1981).
2. A.M. Polyakov, Nucl. Phys. **B268**, 406, (1986).
3. D.I. Gross, T. Piran and S. Weinberg Eds., *Two dimensional quantum gravity and random surfaces*, World Scientific, Singapore, (1992).
4. F. David, P. Ginsparg and Y. Zinn-Justin Eds., *Fluctuating geometries in statistical mechanics and field theory*, Elsevier Science, Amsterdam, (1996).
5. T.L. Curtright, G.I. Chandour, C.B. Thorn and C.K. Zachos, Phys. Rev. Letts, **57**, 799, (1986).
6. T.L. Curtright, G.I. Chandour and C.K. Zachos, Phys. Rev. **D34**, 3811, (1986).
7. E.Braaten, R.D. Pisarski and Sze-Man Tse, Phys. Rev. Letts. **58**, 93, (1987).
8. E. Braaten and C.K. Zachos, Phys. Rev **D35**, 1512, (1987).

9. R. Pisarski, Phys. Rev. Letts. **58**, 1300, (1987).

10. P. Olesen and Sung-Kil Yang, Nucl. Phys. **B283**, 73, (1987).

11. F. David and E. Guitter, Nucl. Phys. **B295**, 332, (1988).

12. B.G.Konopelchenko, Stud. Appl. Math. **96**, 9, (1996).

13. I.A. Taimanov, preprint dg-ga/9511005.

14. K. Kenmotsu, Math. Ann. **245**, 89, (1979).

15. R. Carroll and B.G. Konopelchenko, Int. J. Mod. Phys. **A11**, 1183, (1996).

16. T.I. Wilmore, *Riemannian Geometry*, Clatendon Press, Oxford (1993)

17. Ou-Yang Zhong-can and W. Helfrich, Phys. Rev. **A39**, 5280, (1989).

18. B.A. Dubrovin, S.P. Novikov and A.T. Fomenko, *Modern Geometry*, Nauka Moscow (1979).

19. R. Rajaraman, *Solitons and Istantons*, North-Holland P.C., Amsterdam (1982).

20. A.R. Forsyth, *Theory of differential equations*, part IV, v VI, Dover I. (1906).

21. H. Bateman, *Higher transcendental functions*, v.3, Mc Graw-Hill B.C., New York (1955).

22. A.P. Veselov and S.P. Novikov, Usp. Mat. Nauk, **50**, n. 6,171 (1995)

23. M.A. Olshanetsky and A.M. Perelomov, Phys. Rep., **71**, 314, (1981).

24. B. Sutherland, Phys. Rev., **A5**, 1372, (1972).

25. B.G. Konopelchenko, *Introduction to multidimensional integrable equations*, Plenum Press, New-York (1992).

26. I. A. Taimanov, *Surfaces of revolution in terms of solitons*, dg-ga/9610012.